

PCT


WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification <sup>7</sup> : <b>C07D 215/22, 277/64, A61K 31/428, 31/4375, A61P 7/00, 11/00, 29/00, 35/00</b>		<b>A1</b>	(11) International Publication Number: <b>WO 00/69827</b>
			(43) International Publication Date: 23 November 2000 (23.11.00)
(21) International Application Number: <b>PCT/GB00/01810</b>		(74) Agent: <b>GILL JENNINGS &amp; EVERY</b> ; Broadgate House, 7 Eldon Street, London EC2M 7LH (GB).	
(22) International Filing Date: 12 May 2000 (12.05.00)			
(30) Priority Data: 9911071.0 12 May 1999 (12.05.99) <b>GB</b>		(81) Designated States: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).	
(71) Applicant (for all designated States except US): <b>DARWIN DISCOVERY LIMITED [GB/GB]; Cambridge Science Park, Milton Road, Cambridge CB4 4WE (GB).</b>			
(72) Inventors; and			
(75) Inventors/Applicants (for US only): <b>BAXTER, Andrew, Douglas [GB/GB]; Darwin Discovery Limited, Cambridge Science Park, Milton Road, Cambridge CB4 4WE (GB). OWEN, David, Alan [GB/GB]; Darwin Discovery Limited, Cambridge Science Park, Milton Road, Cambridge CB4 4WE (GB). WATSON, Robert, John [GB/GB]; Darwin Discovery Limited, Cambridge Science Park, Milton Road, Cambridge CB4 4WE (GB). HANNAH, Duncan [GB/GB]; Darwin Discovery Limited, Cambridge Science Park, Milton Road, Cambridge CB4 4WE (GB). BATTY, Duncan [GB/GB]; Darwin Discovery Limited, Cambridge Science Park, Milton Road, Cambridge CB4 4WE (GB).</b>		<b>Published</b> With international search report.	

(54) Title: **HYDROXAMIC AND CARBOXYLIC ACID DERIVATIVES**



(57) Abstract

Compounds of formula (I) in which A is C=O, NR<sub>6</sub>, O, S or a bond; B is a carbon or nitrogen atom; X is C=O, NR<sub>6</sub>, O, S, or a bond; Y is OH or NHOH; and the other variables are as defined in the claims, have therapeutic utility as inhibitors of metalloproteinases.

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

## HYDROXAMIC AND CARBOXYLIC ACID DERIVATIVES

### Field of the Invention

This invention relates to hydroxamic and carboxylic acid derivatives, and to their use in medicine.

### 5 Background to the Invention

Metalloproteinases, including matrix metalloproteinase (MMP), (human fibroblast) collagenase, gelatinase and TNF $\alpha$  convertase (TACE), and their modes of action, and also inhibitors thereof and their clinical effects, are described in WO-A-9611209, WO-A-9712902 and WO-A-9719075, the contents of which are incorporated herein by reference.


- 10 MMP inhibitors may also be useful in the inhibition of other mammalian metalloproteinases such as the ADAM or ADAM-TS families. Members of the ADAM family include TNF $\alpha$  convertase (TACE) and ADAM-10, which can cause the release of TNF $\alpha$  from cells, and others, which have been demonstrated to be expressed by human articular cartilage cells and also involved in the destruction of myelin basic protein, a phenomenon associated with
- 15 multiple sclerosis.

Compounds which have the property of inhibiting the action of metalloproteinases involved in connective tissue breakdown, such as collagenase, stromelysin and gelatinase, have been shown to inhibit the release of TNF $\alpha$  both *in vitro* and *in vivo*. See Gearing *et al* (1994), Nature 370:555-557; McGeehan *et al* (1994), Nature 370:558-561; GB-A-2268934; and WO-A-9320047. All of these reported inhibitors contain a hydroxamic acid zinc-binding group, as do the imidazole-substituted compounds disclosed in WO-A-9523790. Other compounds that inhibit MMP and/or TNF $\alpha$  are described in WO-A-9513289, WO-A-9611209, WO-A-96035687, WO-A-96035711, WO-A-96035712 and WO-A-96035714.

### 25 Summary of the Invention

The invention encompasses novel compounds of formula (I) which are useful inhibitors of matrix metalloproteinases, ADAM or ADAM-TS enzymes, and which are useful for the treatment of disease mediated by those enzymes and/or TNF $\alpha$  mediated diseases, including degenerative diseases and certain cancers.

- 30 Novel compounds according to the invention are of the general type represented by formula (I):



5

wherein

$n = 0-3$ ;

Y is OH or NHOH;

$R^1$  is H,  $R^x$  or a group (optionally substituted with  $R^x$ ) selected from  $C_{1-6}$  alkyl,  $C_{2-6}$  alkenyl,  $C_{2-6}$  alkynyl, aryl,  $C_{1-6}$  alkyl-aryl, heteroaryl,  $C_{1-6}$  alkyl-heteroaryl, cycloalkyl,  $C_{1-6}$  alkyl-cycloalkyl, heterocycloalkyl,  $C_{1-6}$  alkyl-heterocycloalkyl, cycloalkenyl,  $C_{1-6}$  alkyl-cycloalkenyl, heterocycloalkenyl and  $C_{1-6}$  alkyl-heterocycloalkenyl;

$R^2$  is H or  $C_{1-6}$  alkyl;

or  $CR^1R^2$  is cycloalkyl or heterocycloalkyl optionally substituted with  $R^x$ ;

$R^x$  is  $R^3$  or a group (optionally substituted with  $R^3$ ) selected from  $C_{1-6}$  alkyl, aryl,  $C_{1-6}$  alkyl-aryl, heteroaryl and  $C_{1-6}$  alkyl-heteroaryl;

$R^3$  is  $OR^9$ ,  $OCF_3$ ,  $OCF_2H$ ,  $OCH_2F$ ,  $COR^9$ ,  $CO_2R^4$ ,  $CON(R^9)_2$ ,  $N(R^9)_2$ ,  $NR^9COR^9$ ,  $NR^9CON(R^9)_2$ ,  $NR^9CO_2R^{10}$ ,  $NR^9SO_2R^{10}$ ,  $S(O)_{0-2}R^{10}$ ,  $SO_2N(R^9)_2$ , cycloimidyl (optionally substituted with  $R^5$ ) or, where  $R^3$  is not attached to aryl or heteroaryl,  $R^3$  may additionally be  $=O$ ,  $=NOH$  or  $=NOR^{10}$ ;

$R^4$  is H or  $C_{1-6}$  alkyl;

$R^5$  is  $C_{1-6}$  alkyl;

A is  $C=O$ ,  $NR^6$ , O, S, or a bond;

--- represents a single bond and B is  $CHR^6$  or NH, or --- represents a double bond

and B is  $CR^6$  or N;

X is  $C=O$ ,  $NR^6$ , O or S;

provided that when B is  $CR^6$  or  $CHR^6$  and A is a bond, X is not O or S;

$R^6$  is H or a group selected from  $C_{1-6}$  alkyl, aryl,  $C_{1-6}$  alkyl-aryl, heteroaryl and  $C_{1-6}$  alkyl-heteroaryl;

W is an aryl or heteroaryl ring optionally substituted with  $R^7$ ;

$R^7$  is  $R^8$  or a group (optionally substituted with  $R^8$ ) selected from  $C_{1-6}$  alkyl, aryl,  $C_{1-6}$  alkyl-aryl, heteroaryl,  $C_{1-6}$  alkyl-heteroaryl, cycloalkyl,  $C_{1-6}$  alkyl-cycloalkyl, heterocycloalkyl and  $C_{1-6}$  alkyl-heterocycloalkyl;

$R^8$  is selected from  $C=NOR^9$ ,  $N(R^9)_2$ ,  $NR^9COR^9$ ,  $NR^9CON(R^9)_2$ ,  $NR^9CO_2R^{10}$ ,  
 5  $NR^9SO_2R^{10}$ ,  $OR^9$ ,  $OCF_3$ ,  $OCF_2H$ ,  $OCH_2F$ ,  $COR^9$ ,  $CO_2R^4$ ,  $CON(R^9)_2$ ,  $S(O)_{0-2}R^{10}$  and  $SO_2N(R^9)_2$ ;

$R^9$  is H or a group selected from  $C_{1-6}$  alkyl, aryl,  $C_{1-6}$  alkyl-aryl, heteroaryl,  $C_{1-6}$  alkyl-heteroaryl, cycloalkyl,  $C_{1-6}$  alkyl-cycloalkyl, heterocycloalkyl and  $C_{1-6}$  alkyl-heterocycloalkyl, wherein said group is optionally substituted with  $R^{10}$ ,  $COR^{10}$ ,  $SO_{0-2}R^{10}$ ,  
 10  $CO_2R^{10}$ ,  $OR^{10}$ ,  $OCF_3$ ,  $OCF_2H$ ,  $OCH_2F$ ,  $CONR^4R^{10}$ ,  $NR^4R^{10}$  or  $SO_2NR^4R^{10}$  and for each case of  $N(R^9)_2$  the  $R^9$  groups are the same or different or  $N(R^9)_2$  is heterocycloalkyl optionally substituted with  $R^{10}$ ,  $COR^{10}$ ,  $SO_{0-2}R^{10}$ ,  $CO_2R^{10}$ ,  $OR^{10}$ ,  $CONR^4R^{10}$ ,  $NR^4R^{10}$ , or  $SO_2NR^4R^{10}$ ; and

$R^{10}$  is  $C_{1-6}$  alkyl, aryl,  $C_{1-6}$  alkyl-aryl, heteroaryl or  $C_{1-6}$  alkyl-heteroaryl;  
 15 and the salts, solvates, hydrates, N-oxides, protected amino, protected carboxy and protected hydroxamic acid derivatives thereof.

#### Description of the Invention

Preferred compounds of the invention are those wherein any one or more of the following apply:  $n = 1$ ; Y is NHOH;  $CR^1R^2$  forms the said optionally substituted cycloalkyl  
 20 or heterocycloalkyl ring; A is CO, B is  $CR^6$  and X is  $NR^6$ , O or S; A is CO, B is N and X is  $NR^6$ , O or S; A is CO, B is  $CR^6$  and X is CO; A is  $NR^6$ , O or S, B is  $CR^6$  and X is CO; A is CO, B is  $CHR^6$  and X is  $NR^6$ , O or S; A is  $NR^6$ , O or S, B is  $CHR^6$  and X is CO; A is a bond, B is N and X is  $NR^6$ , O or S; A is a bond, B is  $CR^6$  and X is  $NR^6$ , O or S; and A is a bond, B is  $CHR^6$  and X is  $NR^6$ , O or S.

25 It will be appreciated that the compounds according to the invention can contain one or more asymmetrically substituted carbon atoms. The presence of one or more of these asymmetric centres in a compound of formula (I) can give rise to stereoisomers, and in each case the invention is to be understood to extend to all such stereoisomers, including enantiomers and diastereomers, and mixtures including racemic mixtures thereof.  
 30 As used in this specification, alone or in combination, the term " $C_{1-6}$  alkyl" refers to straight or branched chain alkyl moiety having from one to six carbon atoms, including for example, methyl, ethyl, propyl, isopropyl, butyl, *tert*-butyl, pentyl, hexyl and the like.

The term "C<sub>1-8</sub> alkyl" refers to straight or branched chain alkyl moiety having from one to eight carbon atoms, including for example, methyl, ethyl, propyl, isopropyl, butyl, *tert*-butyl, pentyl, hexyl, octyl and the like.

The term "C<sub>2-6</sub> alkenyl" refers to a straight or branched chain alkyl moiety having  
5 two to six carbon atoms and having in addition one double bond, of either E or Z stereochemistry where applicable. This term would include for example, vinyl, 1-propenyl, 1- and 2- butenyl, 2- methyl-2-propenyl etc.

The term "C<sub>2-6</sub> alkynyl" refers to a straight or branched chain alkyl moiety having two to six carbon atoms and having in addition one triple bond. This term would include  
10 for example, ethynyl, 1-propynyl, 1- and 2- butynyl, 1- methyl-2-butynyl etc.

The term "cycloalkyl" refers to a saturated alicyclic moiety having from three to six carbon atoms and includes for example cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl and the like.

The term "cycloalkenyl" refers to an alicyclic moiety having from three to six  
15 carbon atoms and having in addition one double bond. This term includes, for example, cyclopentenyl and cyclohexenyl.

The term "heterocycloalkyl" refers to a saturated heterocyclic moiety having from two to six carbon atoms and one or more heteroatom from the group N, O, S (or oxidised versions thereof) which may be optionally benzofused at any available position. This  
20 includes for example azetidiny, pyrrolidinyl, tetrahydrofuranyl, piperidinyl, benzodioxole and the like.

The term "heterocycloalkenyl" refers to an alicyclic moiety having from three to six carbon atoms and one or more heteroatoms from the group N, O, S and having in addition one double bond. This term includes, for example, dihydropyranyl.

25 The term "aryl" refers to an aromatic carbocyclic radical having a single ring or two condensed rings, optionally substituted with an aryl group substituent. This term includes, for example phenyl or naphthyl.

The term "heteroaryl" refers to aromatic ring systems of five to ten atoms of which at least one atom is selected from O, N and S, and optionally substituted with an aryl  
30 group substituent. This term includes for example furanyl, thiophenyl, pyridyl, indolyl, quinolyl and the like.

The term "aryl group substituent" refers to a substituent chosen from halogen, CN, CF<sub>3</sub>, CHF<sub>2</sub>, CH<sub>2</sub>F, and NO<sub>2</sub>.

The term "halogen" means fluorine, chlorine, bromine or iodine.

The term "benzofused" refers to the addition of a benzene ring sharing a common  
5 bond with the defined ring system.

The term "cycloimidyl" refers to a saturated ring of five to ten atoms containing the atom sequence -C(=O)NC(=O)-. The ring may be optionally benzofused at any available position. Examples include succinimidoyl, phthalimidoyl and hydantoinyl.

The term "optionally substituted" means optionally substituted with one or more  
10 of the groups specified, at any available position or positions.

The terms "protected amino", "protected carboxy" and "protected hydroxamic acid" mean amino, carboxy and hydroxamic acid groups which can be protected in a manner familiar to those skilled in the art. For example, an amino group can be protected by a benzyloxycarbonyl, *tert*-butoxycarbonyl, acetyl or like group, or may be in the form  
15 of a phthalimido or like group. A carboxyl group can be protected in the form of a readily-cleavable ester such as the methyl, ethyl, benzyl or *tert*-butyl ester. A hydroxamic acid may be protected as either N or O-substituted derivatives, such as O-benzyl or O-*tert*-butyldimethylsilyl.

Salts of compounds of formula (I) include pharmaceutically-acceptable salts, for  
20 example acid addition salts derived from inorganic or organic acids, such as hydrochlorides, hydrobromides, p-toluenesulphonates, phosphates, sulphates, perchlorates, acetates, trifluoroacetates, propionates, citrates, malonates, succinates, lactates, oxalates, tartrates and benzoates.

Salts may also be formed with bases. Such salts include salts derived from  
25 inorganic or organic bases, for example alkali metal salts such as magnesium or calcium salts, and organic amine salts such as morpholine, piperidine, dimethylamine or diethylamine salts.

When the "protected carboxy" group in compounds of the invention is an esterified carboxyl group, it may be a metabolically-labile ester of formula CO<sub>2</sub>R<sup>11</sup> where R<sup>9</sup> may be  
30 an ethyl, benzyl, phenethyl, phenylpropyl,  $\alpha$  or  $\beta$ -naphthyl, 2,4-dimethylphenyl, 4-*tert*-butylphenyl, 2,2,2-trifluoroethyl, 1-(benzyloxy)benzyl, 1-(benzyloxy)ethyl, 2-methyl-1-propionyloxypentyl, 2,4,6-trimethylbenzyloxymethyl or pivaloxymethyl group.


Compounds of the general formula (I) may be prepared by any suitable method known in the art and/or by the following processes.

It will be appreciated that, where a particular stereoisomer of formula (I) is required, the synthetic processes described herein may be used with the appropriate homochiral starting material and/or isomers maybe resolved from mixtures using conventional separation techniques (e.g. HPLC).

The compounds according to the invention may be prepared by the following process. In the description and formulae below the groups  $R^1$ ,  $R^2$ ,  $R^3$ ,  $R^4$ ,  $R^5$ ,  $R^6$ ,  $R^7$ ,  $R^8$ ,  $R^9$ ,  $R^{10}$ ,  $R^{11}$ , W, A, B, X and Y are as defined above, except where otherwise indicated.

It will be appreciated that functional groups, such as amino, hydroxyl or carboxyl groups, present in the various compounds described below, and which it is desired to retain, may need to be in protected form before any reaction is initiated. In such instances, removal of the protecting group may be the final step in a particular reaction. Suitable protecting groups for such functionality will be apparent to those skilled in the art. For specific details see Greene *et al*, "Protective Groups in Organic Synthesis", Wiley Interscience.

A process for preparing compounds of general formula (I) comprises reaction of a compound of formula  $CHR^1R^2COY$  (II) with a compound of formula (III) where Z is an appropriate leaving group such as a halogen (for example bromide) or an alkylsulfonate such as methanesulfonate.



25 Suitable conditions for this reaction include the presence of a strong base such as lithium diisopropylamide in an inert solvent such as tetrahydrofuran.

Many compounds of formula (II) are available commercially, or may be prepared from materials available commercially using methods known to those skilled in the art. Compounds of formula (II) may be prepared alternatively from a compound of formula  $R^{12}O_2CCH_2CO_2R^{12}$  (IV) (where  $R^{12}$  is an appropriate protecting group such as ethyl) in a four-step sequence involving (a) alkylation of (IV) with  $R^1Z$  in the presence of a base such as sodium alkoxide to give  $RO_2CCHR^1CO_2R$  (V), (b) alkylation of (V) with  $R^2Z$  in



the presence of an appropriate base such as alkoxide to give  $\text{RO}_2\text{CCR}^1\text{R}^2\text{CO}_2\text{R}$  (VI), (c) conversion of (VI) to the di-acid  $\text{HO}_2\text{CCR}^1\text{CR}^2\text{CO}_2\text{H}$  (VII) by treatment with (where  $\text{R}^{12}$  is ethyl) strong acid such as hydrochloric acid or strong base such as sodium hydroxide, and (d) decarboxylation of (VII) by, for example, the action of heat in the presence of an appropriate catalyst (such as toxic acid) to give (II), where Y is OH. The order of steps (a) and (b) may be reversed, if this is appropriate.

Many compounds of formula (III) are available commercially or may be prepared from compounds available commercially by methods known to those skilled in the art. Compounds of formula (I) may also be prepared by interconversion of other compounds of formula (I). Hydroxamic acids ( $\text{Y}=\text{NHOH}$ ) of general formula (I) may be prepared from carboxylic acids ( $\text{Y}=\text{OH}$ ) of formula (I) or protected versions thereof (such as esters) using methods known to those skilled in the art. Likewise, a compound of formula (I) where  $\text{R}^2$  is not H may be prepared from a compound of formula (VI) where  $\text{R}^2$  is H by reaction with a compound  $\text{R}^2\text{Z}$  (where Z is as defined above) in the presence of a strong base such as lithiumdiisopropylamide in an inert solvent such as tetrahydrofuran. Similarly, intermediates of any appropriate formula may be prepared by the interconversion of other compounds of the same formula.

Any mixtures of final products or intermediates obtained can be separated on the basis of the physico-chemical differences of the constituents, in known manner, into the pure final products or intermediates, for example by chromatography, distillation, fractional crystallization, or by formation of a salt if appropriate or possible under the circumstances.

The compounds according to the invention exhibit *in vitro* inhibiting activities with respect to the stromelysin, collagenase, gelatinase, ADAM or ADAM-TS enzymes. Compounds according to the invention may also exhibit *in vitro* inhibition of membrane shedding events known to be mediated by metalloproteinases, for example,  $\alpha$ -APP, ACE, TGF- $\alpha$ , TNF- $\alpha$ , Fas ligand, selectins, TNFR-I, TNFR-II, CD30, IL-6R, CD43, CD44, CD16-I, CD16-II, Folate receptor, CD23, or IL-1RII.

The activity and selectivity of the compounds may be determined by use of the appropriate enzyme inhibition test, for example as described in Examples A-M of WO-A-98/05635, by the assay for the inhibition of CD23 shedding described in WO-A-99/24399, or by the following assay of TNF RI shedding.

The potency of the compounds of general formula (I) to act as inhibitors of the production of TNF RI is determined using the following procedure. A 100 $\mu$ M solution of the inhibitor being tested or dilutions thereof is incubated at 37° C in an atmosphere of 5% CO<sub>2</sub> with peripheral blood mononuclear cells (PBMC). PBMC are isolated from buffy coats by standard procedures using Ficoll. A 100 $\mu$ M solution of the inhibitor being tested or dilutions thereof is incubated for 22 hours at 37° C in an atmosphere of 5% CO<sub>2</sub> with 1 x 10<sup>6</sup>/ml PBMC stimulated with LPS. The cells are centrifuged down and the supernatant is assayed for TNF RI using a commercially available ELISA kit (@ & D Systems). The activity in the presence of 0.1mM inhibitor or dilutions thereof is compared to activity in a control devoid of inhibitor and results reported as that inhibitor concentration effecting 50% inhibition of the production of TNF RI.

This invention also relates to a method of treatment for patients (including man and/or mammalian animals raised in the dairy, meat or fur industries or as pets) suffering from disorders or diseases which can be attributed to stromelysin as previously described, and more specifically, a method of treatment involving the administration of the matrix metalloproteinase inhibitors of formula (I) as the active constituents.

Accordingly, the compounds of formula (I) can be used among other things in the treatment of osteoarthritis and rheumatoid arthritis, and in diseases and indications resulting from the over-expression of these matrix metalloproteinases such as found in certain metastatic tumour cell lines.

As mentioned above, compounds of formula (I) are useful in human or veterinary medicine since they are active as inhibitors of TNF and MMPs. Accordingly in another aspect, this invention concerns:

a method of management (by which is meant treatment or prophylaxis) of disease or conditions mediated by TNF and/or MMPs in mammals, in particular in humans, which method comprises administering to the mammal an effective, amount of a compound of formula (I) above, or a pharmaceutically acceptable salt thereof; and

a compound of formula (I) for use in human or veterinary medicine, particularly in the management (by which is meant treatment or prophylaxis) of diseases or conditions mediated by TNF and/or MMPs; and

the use of a compound of formula (I) in the preparation of an agent for the management (by which is meant treatment or prophylaxis) of diseases or conditions mediated by TNF and/or MMPs.

The disease or conditions referred to above include inflammatory diseases, autoimmune diseases, cancer, cardiovascular diseases, diseases involving tissue breakdown such as rheumatoid arthritis, osteoarthritis, osteoporosis, neurodegeneration, Alzheimer's disease, stroke, vasculitis, Crohn's disease, ulcerative colitis, multiple sclerosis, periodontitis, gingivitis and those involving tissue breakdown such as bone resorption, haemorrhage, coagulation, acute phase response, cachexia and anorexia, acute infections, bacterial infections, HIV infections, fever, shock states, graft versus host reactions, dermatological conditions, surgical wound healing, psoriasis, atopic dermatitis, epidermolysis bullosa, tumour growth, angiogenesis and invasion by secondary metastases, ophthalmological disease, retinopathy, corneal ulceration, reperfusion injury, migraine, meningitis, asthma, rhinitis, allergic conjunctivitis, eczema, anaphylaxis, restenosis, congestive heart failure, endometriosis, atherosclerosis, endosclerosis, aspirin-independent anti-thrombosis, systemic lupus erythematosus and solid organ transplant.

Compounds of formula (I) may also be useful in the treatment of pelvic inflammatory disease (PID), age-related macular degeneration and cancer-induced bone resorption. Further, they can be used in the treatment of lung diseases, e.g. selected from cystic fibrosis, adult respiratory distress syndrome (ARDS), emphysema, bronchitis obliterans-organising pneumonia (BOOP), idiopathic pulmonary fibrosis (PIF), diffuse alveolar damage, pulmonary Langerhan's cell granulomatosis, pulmonary lymphangioleiomyomatosis (LAM) and chronic obstructive pulmonary disease (COPD).

For the treatment of rheumatoid arthritis, osteoarthritis, and in diseases and indications resulting from the over-expression of matrix metalloendoproteinases such as found in certain metastatic tumour cell lines or other diseases mediated by the matrix metalloendoproteinases or increased TNF production, the compounds of formula (I) may be administered orally, topically, parenterally, by inhalation spray or rectally in dosage unit formulations containing non-toxic pharmaceutically acceptable carriers, adjuvants and vehicles. The term parenteral as used herein includes subcutaneous injections, intravenous, intramuscular, intrasternal injection or infusion techniques. In addition to the treatment

of warm-blooded animals such as mice, rats, horses, cattle, sheep, dogs, cats etc, the compounds of the invention are effective in the treatment of humans.

The pharmaceutical composition containing the active ingredient may be in a form suitable for oral use, for example, as tablets, troches, lozenges, aqueous or oily suspensions, dispersible powders or granules, emulsions, hard or soft capsules, or syrups or elixirs. Compositions intended for oral use may be prepared according to any method known to the art for the manufacture of pharmaceutical compositions and such compositions may contain one or more agents selected from sweetening agents, flavouring agents, colouring agents and preserving agents in order to provide pharmaceutically elegant and palatable preparations. Tablets contain the active ingredient in admixture with non-toxic pharmaceutically acceptable excipients which are suitable for the manufacture of tablets. These excipients may be for example, inert diluents, such as calcium carbonate, sodium carbonate, lactose, calcium phosphate or sodium phosphate; granulating and disintegrating agents, for example corn starch, or alginic acid; binding agents, for example starch, gelatin or acacia, and lubricating agents, for example magnesium stearate, stearic acid or talc. The tablets may be uncoated or they may be coated by known techniques to delay disintegration and absorption in the gastrointestinal tract and thereby provide a sustained action over a longer period. For example, a time delay material such as glyceryl monostearate or glyceryl distearate may be employed. They may also be coated by the techniques described in US-A-4256108, US-A-4166452, and US-A-4265874, to form osmotic therapeutic tablets for control release.

Formulations for oral use may also be presented as hard gelatin capsules where in the active ingredient is mixed with an inert solid diluent, for example calcium carbonate, calcium phosphate or kaolin, or as soft gelatin capsules wherein the active ingredient is mixed with water or an oil medium, for example peanut oil, liquid paraffin or olive oil. Aqueous suspensions contain the active materials in admixture with excipients suitable for the manufacture of aqueous suspensions. Such excipients are suspending agents, for example sodium carboxymethylcellulose, methylcellulose, hydroxy-propylmethylcellulose, sodium alginate polyvinyl-pyrrolidone, gum tragacanth and gum acacia; dispersing or wetting agents may be a naturally occurring phosphatide, for example lecithin, or condensation products of an alkylene oxide with fatty acids, for example polyoxyethylene stearate, or condensation products of ethylene oxide with long chain aliphatic alcohols, for

example heptadecaethyleneoxycetanol, or condensation products of ethylene oxide with partial esters derived from fatty acids and a hexitol such a polyoxyethylene with partial esters derived from fatty acids and hexitol anhydrides, for example polyoxyethylene sorbitan monooleate. The aqueous suspensions may also contain one or more  
5 preservatives, for example ethyl, or n-propyl, p-hydroxybenzoate, one or more colouring agents, one or more flavouring agents, and one or more sweetening agents, such as sucrose or saccharin.

Oily suspensions may be formulated by suspending the active ingredient in a vegetable oil, for example arachis oil, olive oil, sesame oil or coconut oil, or in a mineral  
10 oil such as liquid paraffin. The oily suspensions may contain a thickening agent, for example beeswax, hard paraffin or cetyl alcohol. Sweetening agents such as those set forth above, and flavouring agents may be added to provide a palatable oral preparation. These compositions may be preserved by the addition of an anti-oxidant such as ascorbic acid.

15 Dispersible powders and granules suitable for preparation of an aqueous suspension by the addition of water provide the active ingredient in admixture with a dispersing or wetting agent, suspending agent and one or more preservatives. Suitable dispersing or wetting agents and suspending agents are exemplified, for example sweetening, flavouring and colouring agents may also be present.

20 The pharmaceutical compositions of the invention may also be in the form of oil-in-water emulsions. The oily phase may be a vegetable oil, for example olive oil or arachis oil, or a mineral oil, for example liquid paraffin or mixtures of these. Suitable emulsifying agents may be naturally occurring gums, for example gum acacia or gum tragacanth, naturally occurring phosphatides, for example soya bean, lecithin, and esters or partial  
25 esters derived from fatty acids and hexitol anhydrides, for example sorbitan monooleate and condensation products of the said partial esters with ethylene oxide, for example polyoxyethylene sorbitan monooleate. The emulsions may also contain sweetening and flavouring agents.

Syrups and elixirs may be formulated with sweetening agents, for example glycerol,  
30 propylene glycol, sorbitol or sucrose. Such formulations may also contain a demulcent, a preservative and flavouring and colouring agents. The pharmaceutical compositions may be in the form of a sterile injectable aqueous or oleagenous suspension. This suspension

may be formulated according to the known art using those suitable dispersing or wetting agents and suspending agents which have been mentioned above. The sterile injectable preparation may also be in a sterile injectable solution or suspension in a non-toxic parenterally acceptable diluent or solvent, for example as a solution in 1,3-butanediol.

- 5 Among the acceptable vehicles and solvents that may be employed are water, Ringer's solution and isotonic sodium chloride solution. In addition, sterile, fixed oils are conventionally employed as a solvent or suspending medium. For this purpose any bland fixed oil may be employed including synthetic mono- or diglycerides. In addition, fatty acids such as oleic acid find use in the preparation of injectables.

- 10 The compounds of formula (I) may also be administered in the form of suppositories for rectal administration of the drug. These compositions can be prepared by mixing the drug with a suitable non-irritating excipient which is solid at ordinary temperatures but liquid at the rectal temperature and will therefore melt in the rectum to release the drug. Such materials are cocoa butter and polyethylene glycols.

- 15 For topical use, creams, ointments, jellies, solutions or suspensions, etc containing the compounds of Formula (I) are employed. For the purposes of this specification, topical application includes mouthwashes and gargles.

- Dosage levels of the order of from about 0.05 mg to about 140 mg per kilogram of body weight per day are useful in the treatment of the above- indicated conditions (about 2.5 mg to about 7 g per patient per day). For example, inflammation may be effectively treated by the administration of from about 0.01 to 50 mg of the compound per kilogram of body weight per day (about 0.5 mg to about 3.5 g per patient per day).

- The amount of active ingredient that may be combined with the carrier materials to produce a single dosage form will vary depending upon the host treated and the particular mode of administration. For example, a formulation intended for the oral administration of humans may vary from about 5 to about 95% of the total composition. Dosage unit forms will generally contain between from about 1 mg to about 500 mg of active ingredient.

- It will be understood, however, that the specific dose level for any particular patient will depend upon a variety of factors including the activity of the specific compound employed, the age, body weight, general health, sex, diet time of

administration, route of administration, rate of excretion, drug combination and the severity of the particular disease undergoing therapy.

The following Examples illustrate the invention.

**Example 1    2-(4-Oxo-6-trifluoromethoxy-1,4-dihydroquinolin-2-yl)acetic acid N-hydroxyamide**


To a suspension of (4-oxo-6-trifluoromethoxy-1,4-dihydroquinolin-2-yl)acetic acid ethyl ester (0.020 g) in ethanol (1 ml) at room temperature was added aqueous hydroxylamine (0.1 ml). The mixture was left stirring at room temperature for 48 hours. The resultant white suspension was diluted with water (10 ml), filtered, washed with water (15 ml) and dried *in vacuo* to provide the title compound as a white solid (0.007 g, 39%).  
 $R_f=0.18$  (10% methanol/dichloromethane)  
MS=303 (M+H)

**Example 2    3-Benzothiazol-2-ylpropionic acid N-hydroxyamide**

To a solution of 3-benzothiazol-2-yl-propionic acid (0.060 g) in dichloromethane (5 ml) at 0°C was added *O*-(*tert*-butyldimethylsilyl)hydroxylamine (0.046 g) and 3-ethyl-1-(3-dimethylaminopropyl)carbodiimide (0.072 g). The reaction mixture was stirred for 3 hours and allowed to warm to room temperature. The solvent was evaporated *in vacuo*. The residue was then dissolved in ethyl acetate (25 ml), washed with water (15 ml), saturated aqueous sodium hydrogen carbonate solution (2 x 15 ml), water (2 x 15 ml) and saturated brine (20 ml), dried over anhydrous sodium sulfate and evaporated under reduced pressure. The resultant gum was dissolved in tetrahydrofuran (15 ml), cooled in ice and treated with *tert*-butylammonium fluoride, as a 1.0 M solution in tetrahydrofuran (0.3 ml). The tetrahydrofuran was evaporated and ethyl acetate (20 ml) was added. This solution was then washed with water (15 ml), saturated aqueous sodium hydrogen carbonate solution (2 x 15 ml) and water (15 ml). The combined aqueous washes were back-extracted with ethyl acetate (2 x 15 ml) and the combined organic extracts were washed with saturated brine (20 ml), dried over anhydrous sodium sulfate and evaporated under reduced pressure. The residue was then triturated with hexane (10 ml) and extracted into diethyl ether (2 x 10 ml). The ether was then evaporated and residue was purified by flash column chromatography with 8% methanol/dichloromethane as eluent, to yield the title compound as a white solid (0.010 g, 16%).  
 $R_f=0.43$  (10% methanol/dichloromethane)  
MS=223 (M+H)

CLAIMS

1. Use of a compound for the manufacture of a medicament for the treatment or prevention of a condition associated with matrix metalloproteinases or that is mediated by ADAM or ADAM-TS enzymes, a condition that is mediated by TNF $\alpha$ , or a condition that is mediated by a metalloproteinase, wherein the compound is of formula (I)



wherein  $n = 0-3$ ;

Y is OH or NHOH;

- 15  $R^1$  is H,  $R^2$  or a group (optionally substituted with  $R^2$ ) selected from  $C_{1-6}$  alkyl,  $C_{2-6}$  alkenyl,  $C_{2-6}$  alkynyl, aryl,  $C_{1-6}$  alkyl-aryl, heteroaryl,  $C_{1-6}$  alkyl-heteroaryl, cycloalkyl,  $C_{1-6}$  alkyl-cycloalkyl, heterocycloalkyl,  $C_{1-6}$  alkyl-heterocycloalkyl, cycloalkenyl,  $C_{1-6}$  alkyl-cycloalkenyl, heterocycloalkenyl and  $C_{1-6}$  alkyl-heterocycloalkenyl;

$R^2$  is H or  $C_{1-6}$  alkyl;

- 20 or  $CR^1R^2$  is cycloalkyl or heterocycloalkyl optionally substituted with  $R^2$ ;

$R^2$  is  $R^3$  or a group (optionally substituted with  $R^3$ ) selected from  $C_{1-6}$  alkyl, aryl,  $C_{1-6}$  alkyl-aryl, heteroaryl and  $C_{1-6}$  alkyl-heteroaryl;

- $R^3$  is  $OR^9$ ,  $OCF_3$ ,  $OCF_2H$ ,  $OCH_2F$ ,  $COR^9$ ,  $CO_2R^4$ ,  $CON(R^9)_2$ ,  $N(R^9)_2$ ,  $NR^9COR^9$ ,  $NR^9CON(R^9)_2$ ,  $NR^9CO_2R^{10}$ ,  $NR^9SO_2R^{10}$ ,  $S(O)_{0-2}R^{10}$ ,  $SO_2N(R^9)_2$ , cycloimidyl (optionally substituted with  $R^5$ ) or, where  $R^3$  is not attached to aryl or heteroaryl,  $R^3$  may additionally be  $=O$ ,  $=NOH$  or  $=NOR^{10}$ ;
- 25

$R^4$  is H or  $C_{1-6}$  alkyl;

$R^5$  is  $C_{1-6}$  alkyl;

A is  $C=O$ ,  $NR^6$ , O, S, or a bond;

- 30  $---$  represents a single bond and B is  $CHR^6$  or NH, or  $---$  represents a double bond and B is  $CR^6$  or N;

X is  $C=O$ ,  $NR^6$ , O or S;



provided that when B is CR<sup>6</sup> or CHR<sup>6</sup> and A is a bond, X is not O or S;

W is aryl or heteroaryl optionally substituted with R<sup>7</sup>;

R<sup>7</sup> is H, R<sup>8</sup> or a group (optionally substituted with R<sup>8</sup>) selected from C<sub>1-6</sub> alkyl, aryl, C<sub>1-6</sub> alkyl-aryl, heteroaryl, C<sub>1-6</sub> alkyl-heteroaryl, cycloalkyl, C<sub>1-6</sub> alkyl-cycloalkyl, heterocycloalkyl and C<sub>1-6</sub> alkyl-heterocycloalkyl;

R<sup>8</sup> is selected from C=NOR<sup>9</sup>, N(R<sup>9</sup>)<sub>2</sub>, NR<sup>9</sup>COR<sup>9</sup>, NR<sup>9</sup>CON(R<sup>9</sup>)<sub>2</sub>, NR<sup>9</sup>CO<sub>2</sub>R<sup>10</sup>, NR<sup>9</sup>SO<sub>2</sub>R<sup>10</sup>, OR<sup>9</sup>, OCF<sub>3</sub>, OCF<sub>2</sub>H, OCH<sub>2</sub>F, COR<sup>9</sup>, CO<sub>2</sub>R<sup>4</sup>, CON(R<sup>9</sup>)<sub>2</sub>, S(O)<sub>0-2</sub>R<sup>10</sup> and SO<sub>2</sub>N(R<sup>9</sup>)<sub>2</sub>; and

R<sup>9</sup> is H or a group selected from C<sub>1-6</sub> alkyl, aryl, C<sub>1-6</sub> alkyl-aryl, heteroaryl, C<sub>1-6</sub> alkyl-heteroaryl, cycloalkyl, C<sub>1-6</sub> alkyl-cycloalkyl, heterocycloalkyl and C<sub>1-6</sub> alkyl-heterocycloalkyl, wherein said group is optionally substituted with R<sup>10</sup>, COR<sup>10</sup>, SO<sub>0-2</sub>R<sup>10</sup>, CO<sub>2</sub>R<sup>10</sup>, OR<sup>10</sup>, OCF<sub>3</sub>, OCF<sub>2</sub>H, OCH<sub>2</sub>F, CONR<sup>4</sup>R<sup>10</sup>, NR<sup>4</sup>R<sup>10</sup>, or SO<sub>2</sub>NR<sup>4</sup>R<sup>10</sup>, and for each case of N(R<sup>9</sup>)<sub>2</sub> the R<sup>9</sup> groups are the same or different, or N(R<sup>9</sup>)<sub>2</sub> is heterocycloalkyl optionally substituted with R<sup>10</sup>, COR<sup>10</sup>, SO<sub>0-2</sub>R<sup>10</sup>, CO<sub>2</sub>R<sup>10</sup>, OR<sup>10</sup>, CONR<sup>4</sup>R<sup>10</sup>, NR<sup>4</sup>R<sup>10</sup> or SO<sub>2</sub>NR<sup>4</sup>R<sup>10</sup>;

or a salt, solvate, hydrate, N-oxide, protected amino, protected carboxy or protected hydroxamic acid derivative thereof.

2. Use according to claim 1, wherein CR<sup>1</sup>R<sup>2</sup> forms the said optionally substituted cycloalkyl or heterocycloalkyl ring.

3. Use according to claim 1 or claim 2, wherein A is CO, B is CR<sup>6</sup> and X is NR<sup>6</sup>, O or S; A is CO, B is N and X is NR<sup>6</sup>, O or S; A is CO, B is CR<sup>6</sup> and X is CO; A is NR<sup>6</sup>, O or S, B is CR<sup>6</sup> and X is CO; A is CO, B is CHR<sup>6</sup> and X is NR<sup>6</sup>, O or S; A is NR<sup>6</sup>, O or S, B is CHR<sup>6</sup> and X is CO; A is a bond, B is N and X is NR<sup>6</sup>, O or S; A is a bond, B is CR<sup>6</sup> and X is NR<sup>6</sup>, O or S; and A is a bond, B is CHR<sup>6</sup> and X is NR<sup>6</sup>, O or S.

4. Use according to any preceding claim, wherein n is 1.

5. Use according to any preceding claim, wherein the compound is chiral and in the form of a single enantiomer or diastereomer.

6. Use according to any preceding claim, wherein

n = 0-1;

R<sup>1</sup> is H, R<sup>3</sup> or a group (optionally substituted with R<sup>3</sup>) selected from C<sub>1-6</sub> alkyl, C<sub>2-6</sub> alkenyl, C<sub>2-6</sub> alkynyl, aryl, C<sub>1-6</sub> alkyl-aryl, heteroaryl, C<sub>1-6</sub> alkyl-heteroaryl, cycloalkyl, C<sub>1-6</sub> alkyl-cycloalkyl, heterocycloalkyl and C<sub>1-6</sub> alkyl-heterocycloalkyl;

$R^2$  is H or  $C_{1-6}$  alkyl;

or  $CR^1R^2$  is cycloalkyl or heterocycloalkyl optionally substituted with  $R^3$  or a group (optionally substituted with  $R^3$ ) selected from  $C_{1-6}$  alkyl, aryl,  $C_{1-6}$  alkyl-aryl, heteroaryl and  $C_{1-6}$  alkyl-heteroaryl;

5  $R^3$  is  $OR^9$ ,  $COR^9$ ,  $CO_2R^4$ ,  $CON(R^9)_2$ ,  $N(R^9)_2$ ,  $NR^9COR^9$ ,  $NR^9CON(R^9)_2$ ,  $NR^9CO_2R^{10}$ ,  $NR^9SO_2R^{10}$ ,  $S(O)_{0-2}R^{10}$ ,  $SO_2N(R^9)_2$  or cycloimidyl (optionally substituted with  $R^3$ ); and

$R^6$  is H or  $C_{1-6}$  alkyl.

7. A compound as defined in any preceding claim, independent of use, wherein Y is  
10 NHOH.

8. A compound of claim 7, which is  
2-(4-oxo-6-trifluoromethoxy-1,4-dihydroquinolin-2-yl)acetic acid N-hydroxyamide or  
3-benzothiazol-2-ylpropionic acid N-hydroxyamide

9. A compound of claim 7, for therapeutic use.

15 10. A pharmaceutical composition for use in therapy, comprising a compound of claim 7 or claim 8, and a pharmaceutically-acceptable diluent or carrier.

11. Use of a compound of claim 7 or claim 8, for the manufacture of a medicament for the treatment or prevention of a condition as defined in claim 1.

12. Use according to any of claims 1 to 6 and 11, wherein the condition is selected  
20 from cancer, inflammation and inflammatory diseases, tissue degeneration, periodontal disease, ophthalmological disease, dermatological disorders, fever, cardiovascular effects, haemorrhage, coagulation and acute phase response, cachexia, anorexia, acute infection, HIV infection, shock states, graft versus host reactions, autoimmune disease, reperfusion injury, meningitis, migraine and aspirin-independent anti-thrombosis.

25 13. Use according to any of claims 1 to 6 and 11, wherein the condition is a bacterial infection.

14. Use according to any of claims 1 to 6 and 11, wherein the condition is selected from tumour growth, angiogenesis, tumour invasion and spread, metastases, malignant ascites and malignant pleural effusion.

30 15. Use according to any of claims 1 to 6 and 11, wherein the condition is selected from cerebral ischaemia, ischaemic heart disease, rheumatoid arthritis, osteoarthritis,

osteoporosis, asthma, multiple sclerosis, neurodegeneration, Alzheimer's, atherosclerosis, stroke and vasculitis.

16. Use according to any of claims 1 to 6 and 11, wherein the condition is selected from Crohn's disease and ulcerative colitis.
- 5 17. Use according to any of claims 1 to 6 and 11, wherein the condition is selected from corneal ulceration, retinopathy and surgical wound healing.
18. Use according to any of claims 1 to 6 and 11, wherein the condition is selected from psoriasis, atopic dermatitis, chronic ulcers and epidermolysis bullosa.
19. Use according to any of claims 1 to 6 and 11, wherein the condition is selected  
10 from periodontitis and gingivitis.
20. Use according to any of claims 1 to 6 and 11, wherein the condition is selected from rhinitis, allergic conjunctivitis, eczema and anaphylaxis.
21. Use according to any of claims 1 to 6 and 11, wherein the condition is selected from restenosis, congestive heart failure, endometriosis, atherosclerosis and endosclerosis.
- 15 22. Use according to any of claims 1 to 6 and 11, wherein the condition is selected from pelvic inflammatory disease (PID) and cancer-induced bone resorption.
23. Use according to any of claims 1 to 6 and 11, wherein the condition is age-related macular degeneration.
24. Use according to any of claims 1 to 6 and 11, wherein the condition is selected  
20 from systemic lupus erythematosus and solid organ transplant.
25. Use according to any of claims 1 to 6 and 11, wherein the condition is a lung disease.
26. Use according to claim 25, wherein the condition is selected from cystic fibrosis adult respiratory distress syndrome (ARDS), emphysema, bronchitis obliterans-organising pneumonia (BOOP), idiopathic pulmonary fibrosis (PIF), diffuse alveolar damage, pulmonary Langerhan's cell granulomatosis, pulmonary lymphangioleiomyomatosis (LAM)  
25 and chronic obstructive pulmonary disease (COPD).

# INTERNATIONAL SEARCH REPORT

International Application No

PCT/GB 00/01810

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07D215/22 C07D277/64 A61K31/428 A61K31/4375 A61P7/00  
A61P11/00 A61P29/00 A61P35/00

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07C C07D A61K A61P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

WPI Data, EPO-Internal, CHEM ABS Data

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P, X	EP 0 950 656 A (SHIONOGI) 20 October 1999 (1999-10-20) page 2 -page 3; example 21	1,7,9-12
X	& WO 97 27174 A 31 July 1997 (1997-07-31)	
X	FR 2 338 041 A (DELALANDE) 12 August 1977 (1977-08-12) page 1; examples 1,2	7,9,10
	-/-	

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

\* Special categories of cited documents :

\*A\* document defining the general state of the art which is not considered to be of particular relevance

\*E\* earlier document but published on or after the international filing date

\*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

\*O\* document referring to an oral disclosure, use, exhibition or other means

\*P\* document published prior to the international filing date but later than the priority date claimed

\*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

\*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

\*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

\*&\* document member of the same patent family

Date of the actual completion of the international search

8 August 2000

Date of mailing of the international search report

31/08/2000

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

English, R

## INTERNATIONAL SEARCH REPORT

Int'l. Jonal Application No

PCT/GB 00/01810

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	M. DESCAMPS, ET AL.: "Recherches dans la série des benzo'b!thiophènes. II. Acides benzo'b!thiényl-2 ou -3 acétiques, agents anti-inflammatoires potentiels" CHIMIE THERAPEUTIQUE, vol. 8, no. 5, September 1973 (1973-09), pages 536-544, XP002132934 Editions Dimeo, Arceuil, FR ISSN: 0009-4374 compound 18	7,9,10
X	GB 1 574 822 A (LABORATOIRE L. LAFON) 10 September 1980 (1980-09-10) example 19	7,9,10
X	GB 1 417 822 A (DELALANDE) 17 December 1975 (1975-12-17) example	7,9,10
X	CHEMICAL ABSTRACTS, vol. 70, no. 9, 3 March 1969 (1969-03-03) Columbus, Ohio, US; abstract no. 37689c, F.S. BABICHEV, ET AL.: "(Benzo-2-thiazoyl)alkane(arene)carboxylic acids and their derivatives. VII. Hydrazides, hydroxamic acids, nitriles, and thioamides from (benzo-2-thiazoyl)alkanecarboxylic acids" page 339; XP002144504 abstract: compounds IV & UKRAINSKII KHIMICHESKII ZHURNAL, vol. 34, no. 9, 1968, pages 933-936,	7
A	WO 98 16503 A (AMERICAN CYANAMID) 23 April 1998 (1998-04-23) the whole document	1

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/GB 00/01810

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
EP 0950656	A	20-10-1999	AU 715764 B	10-02-2000
			AU 1319597 A	20-08-1997
			BR 9707010 A	20-07-1999
			HU 9903687 A	28-03-2000
			NO 983376 A	14-09-1998
			SK 98498 A	13-04-1999
			CA 2242416 A	31-07-1997
			CN 1214041 A	14-04-1999
			CZ 9802252 A	16-12-1998
			WO 9727174 A	31-07-1997
			PL 328270 A	18-01-1999
FR 2338041	A	12-08-1977	NONE	
GB 1574822	A	10-09-1980	AT 356078 B	10-04-1980
			AT 193077 A	15-09-1979
			AT 374191 B	26-03-1984
			AT 501480 A	15-08-1983
			AT 361932 B	10-04-1981
			AT 839878 A	15-09-1980
			AT 358556 B	25-09-1980
			AT 839978 A	15-02-1980
			AT 362793 B	10-06-1981
			AT 840078 A	15-11-1980
			AU 516473 B	04-06-1981
			AU 2334477 A	21-09-1978
			BE 852738 A	22-09-1977
			CA 1120928 A	30-03-1982
			CA 1130301 A	24-08-1982
			CH 620894 A	31-12-1980
			CS 200511 B	15-09-1980
			DD 129645 A	01-02-1978
			DE 2711451 A	06-10-1977
			DK 126677 A	24-09-1977
			ES 457105 A	16-10-1978
			FI 770859 A,B,	24-09-1977
			FI 821213 A,B,	06-04-1982
			FI 821214 A,B,	06-04-1982
			FI 821215 A,B,	06-04-1982
			FR 2345430 A	21-10-1977
			FR 2453148 A	31-10-1980
			FR 2453133 A	31-10-1980
			FR 2453158 A	31-10-1980
			IE 44721 B	10-03-1982
			IL 51705 A	30-09-1982
			JP 1404487 C	09-10-1987
			JP 52144601 A	02-12-1977
			JP 62008424 B	23-02-1987
			KR 8001032 A	26-09-1980
			LU 76989 A	18-07-1977
			NL 7703168 A,C	27-09-1977
			NO 771006 A,B,	26-09-1977
			NO 803336 A,B,	26-09-1977
			NO 803337 A,B,	26-09-1977
			NO 803338 A,B,	26-09-1977
			NZ 183616 A	28-03-1979
			PH 16271 A	25-08-1983
			PT 66300 A,B	01-04-1977

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No.

PCT/GB 00/01810

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
GB 1574822 A		SE 432420 B	02-04-1984
		SE 7703263 A	24-09-1977
		SE 452155 B	16-11-1987
		SE 8302171 A	19-04-1983
		SE 458605 B	17-04-1989
		SE 8302172 A	19-04-1983
GB 1417822 A	17-12-1975	FR 2218092 A	13-09-1974
		AU 476756 B	30-09-1976
		AU 6511074 A	07-08-1975
		BE 810683 A	06-08-1974
		CA 1012542 A	21-06-1977
		CH 578570 A	13-08-1976
		DE 2404413 A	29-08-1974
		ES 423368 A	01-06-1976
		JP 49110700 A	22-10-1974
		LU 69362 A	01-10-1974
		NL 7402222 A	21-08-1974
		SE 382815 B	16-02-1976
		SU 501672 A	30-01-1976
		US 3922280 A	25-11-1975
		ZA 7400404 A	27-11-1974
WO 9816503 A	23-04-1998	AU 5145898 A	11-05-1998
		BR 9712525 A	19-10-1999
		EP 0938471 A	01-09-1999
		ZA 9709233 A	15-04-1999